

Exam in Calculus

First Year at the Faculty of Engineering and Science
and the Technical Faculty of IT and Design

5 January 2018

The present exam set consists of 8 numbered pages with 12 multiple-choice problems. A number of points are specified for each question. The total number of points equals 100.

It is allowed to use books, notes etc. It **is not** allowed to use electronic devices.

Your answers must be marked directly in this exam set. The grading of the exam is based solely on these marks. Remember to write your **full name** and **student number** below.

Good luck!

NAME: _____

STUDENT NUMBER: _____

Answers

Problem 1 (9 points)

(a) (5 points). A second order differential equation is given by

$$y'' + 4y' + 29y = 0.$$

Below is a list of function expressions containing two arbitrary constants c_1 and c_2 . Mark the expression which constitute the general solution of the differential equation.

- $y(t) = c_1e^{3t} + c_2e^{5t}$
- $y(t) = c_1e^{2t} + c_2e^{5t}$
- $y(t) = c_1e^{-2t} + c_2e^{3t}$
- $y(t) = c_1e^{2t} + c_2te^{2t}$
- $y(t) = c_1e^{-3t} + c_2te^{-3t}$
- $y(t) = c_1e^{-2t} \cos(5t) + c_2e^{-2t} \sin(5t)$
- $y(t) = c_1e^{-t} \cos(4t) + c_2e^{-t} \sin(4t)$
- $y(t) = c_1e^{2t} \cos(3t) + c_2e^{2t} \sin(3t)$

(b) (4 points). Now consider the inhomogeneous equation

$$y'' + 4y' + 29y = 13e^{-3t}.$$

Which one of the following functions is a particular solution of that equation?

- | | | | |
|--|---|--|--|
| <input type="checkbox"/> $t^2 + e^{-3t+1}$ | <input type="checkbox"/> $2e^{3t}$ | <input type="checkbox"/> $\frac{1}{3}e^{3t}$ | <input checked="" type="checkbox"/> $\frac{1}{2}e^{-3t}$ |
| <input type="checkbox"/> $2e^{-3t}$ | <input type="checkbox"/> $\frac{1}{4}e^{-3t}$ | <input type="checkbox"/> $\frac{1}{9}e^t + e^{-t}$ | <input type="checkbox"/> $\frac{2}{3}e^{-3t+1}$ |

Problem 2 (6 points)

A surface \mathcal{F} in space is determined by the equation $F(x, y, z) = 0$ where

$$F(x, y, z) = x^3 - y^3 + 2z^3 - 9.$$

The surface \mathcal{F} has a tangent plane at the point $P = (2, 1, 1)$. Mark an equation for this tangent plane among the following options:

$3x - 3y + 12z = 15$

$2x - 2y + 7z = 5$

$x - 2y + z = 1$

$-3x + 2y + 6z = 11$

$9x + y + 6z = 25$

$x - 2y - z = 1$

$12x - 3y + 6z = 27$

$x - 3y + 5z = -3$

Problem 3 (10 points)

A plane curve is given by

$$\begin{aligned}x &= t - 2t^2, \\y &= 2t + t^2,\end{aligned}$$

where the parameter t runs through the real numbers.

(a) (2 points). Which point on the curve corresponds to the parameter value $t = 1$?

$(1, 4)$

$(-1, 1)$

$(-1, 2)$

$(3, 4)$

$(-1, 3)$

$(5, 4)$

$(-3, 4)$

$(-5, 4)$

(b) (4 points). What is the curvature of the curve for $t = 1$?

$\frac{\sqrt{2}}{4}$

$\frac{\sqrt{3}}{4}$

$\frac{2}{25}$

5

$\frac{1}{2}$

$\sqrt{3}$

$\frac{1}{9}$

$\sqrt{11}$

(c) (4 points). For which value of t is the curvature maximal?

-2

0

3

5

$-\frac{1}{2}$

$\frac{3}{2}$

$\frac{7}{2}$

9

Problem 4 (7 points)

A function is defined by

$$f(x) = (4x + 1)^{\frac{1}{2}}.$$

(a) (3 points). What is the double derivative $f''(x)$?

- $(4x + 1)^{-\frac{1}{2}}$ $2x^{-\frac{1}{2}}$ $\frac{1}{4}(4x + 1)^{\frac{1}{2}}$
 $-\frac{1}{4}(4x + 1)^{-\frac{3}{2}}$ $6x^{-\frac{3}{2}}$ $-4(4x + 1)^{-\frac{3}{2}}$

(b) (4 points). Which one of the polynomials below is the third order Taylor polynomial for f at the point $x = 0$?

- $1 + 2x - \frac{1}{2}x^2 + \frac{1}{6}x^3$ $1 - \frac{1}{2}x^3$ $2 + 4x^2 - \frac{1}{2}x^3$
 $1 - x + x^2 - 6x^3$ $1 + 3x + x^2 + 2x^3$ $1 + 2x - 2x^2 + 4x^3$

Problem 5 (10 points)

A particle is moving along a curve in space. The position vector of the particle at time t is

$$\mathbf{r}(t) = \langle e^t, e^{-t}, \sqrt{2}t \rangle.$$

(a) (3 points). What is the velocity vector $\mathbf{v}(t)$?

- $\langle e^t, -e^{-t}, \frac{1}{2\sqrt{2}}t \rangle$ $\langle te^{t-1}, -te^{-t-1}, 0 \rangle$
 $\langle e^t, e^{-t}, 1 \rangle$ $\langle e^t, -e^{-t}, \sqrt{2} \rangle$
 $\langle e^t, -e^{-t}, \frac{\sqrt{2}}{2}t^2 \rangle$ $\langle e^t, e^{-t}, t \rangle$

(b) (4 points). What is the speed $v(t)$?

- $e^t + e^{-t}$ $\sqrt{e^{2t} + e^{-2t} + \frac{1}{8}t}$
 $e^t + e^{-t} + \sqrt{2}$ $2e^{2t} + 2t$
 $\sqrt{e^{2t} + e^{-2t} + t^2}$ $\sqrt{e^{2t} + e^{-2t} + 2t^2}$

(c) (3 points). The particle runs through a segment of the motion curve during the time interval $0 \leq t \leq 1$. What is the arc length of that segment?

- e^2 $e - e^{-1}$
 $2e + 1$ $e - e^{-1} + \sqrt{2}$
 $\sqrt{2}$ $3 \ln(2)$

Problem 6 (6 points)

Two complex numbers are given by

$$z_1 = \frac{1+i}{3-2i}, \quad z_2 = \frac{2i}{1+i}$$

(a) (3 points). What is z_1 written in standard form?

- | | | | |
|---|--|-------------------------------|---|
| <input type="checkbox"/> $2 + 3i$ | <input checked="" type="checkbox"/> $\frac{1}{13} + \frac{5}{13}i$ | <input type="checkbox"/> $8i$ | <input type="checkbox"/> $\frac{1}{4} - \frac{3}{4}i$ |
| <input type="checkbox"/> $\frac{1}{5} - \frac{2}{5}i$ | <input type="checkbox"/> $\frac{2}{5} + \frac{3}{5}i$ | <input type="checkbox"/> -5 | <input type="checkbox"/> $-2 - i$ |

(b) (3 points). What is z_2 written in polar form?

- | | | | |
|--|--|---|---|
| <input type="checkbox"/> $2e^{i\pi/3}$ | <input type="checkbox"/> $2e^{i\pi}$ | <input type="checkbox"/> $2e^{i\pi/2}$ | <input type="checkbox"/> $\sqrt{6}e^{i\pi/4}$ |
| <input checked="" type="checkbox"/> $\sqrt{2}e^{i\pi/4}$ | <input type="checkbox"/> $\sqrt{2}e^{-i\pi/3}$ | <input type="checkbox"/> $\frac{1}{2}e^{-i\pi}$ | <input type="checkbox"/> $5e^{3i\pi/4}$ |

Problem 7 (8 points)

A function is defined as

$$f(x, y) = \frac{4x - 2y^2 - 3}{x^2 - y^2}.$$

Mark the correct option in each question below.

(a) (4 points). The domain of f consists of all those points (x, y) which satisfy

- | | |
|--|--|
| <input type="checkbox"/> $x > 0$ and $y > 0$ | <input checked="" type="checkbox"/> $y \neq x$ and $y \neq -x$ |
| <input type="checkbox"/> $x > y$ | <input type="checkbox"/> $x \neq 0$ and $y \neq 0$ |
| <input type="checkbox"/> $4x + 5 \geq 0$ | <input type="checkbox"/> $x \neq 0$ or $y \neq 0$ |

(b) (4 points). The level curve with equation $f(x, y) = 1$ can be described as:

- A parabola with equation $y = 2x^2 + \frac{5}{2}$.
- A parabola with equation $y = x^2 - 2x + 4$.
- A straight line through $(0, 5)$ with slope 4.
- A straight line through $(0, 2)$ with slope 2.
- A circle with center at $(-2, 1)$ and radius $\frac{1}{2}$.
- A circle with center at $(2, 0)$ and radius 1.

Problem 8 (14 points)

A function is defined as

$$f(x, y) = xy^2 - 6x^2 - 3y^2.$$

(a) (2 points). What is the function value $f(1, 1)$?

- | | | | |
|--|----------------------------|----------------------------|-----------------------------|
| <input type="checkbox"/> -10 | <input type="checkbox"/> 0 | <input type="checkbox"/> 3 | <input type="checkbox"/> 7 |
| <input checked="" type="checkbox"/> -8 | <input type="checkbox"/> 2 | <input type="checkbox"/> 5 | <input type="checkbox"/> 11 |

(b) (4 points). Which of the following points are critical points for f ? (*Remark:* Each wrong mark in this question will cancel a true mark.)

- | | | | |
|--|--|---|-----------------------------------|
| <input checked="" type="checkbox"/> (0, 0) | <input type="checkbox"/> (-1, 1) | <input type="checkbox"/> (2, 3) | <input type="checkbox"/> (4, 2) |
| <input type="checkbox"/> (1, 1) | <input checked="" type="checkbox"/> (3, 6) | <input checked="" type="checkbox"/> (3, -6) | <input type="checkbox"/> (4, -16) |

(c) (4 points). What is the directional derivative $D_{\mathbf{u}}f(P)$ at the point $P = (-1, 1)$ and in the direction of the unit vector

$$\mathbf{u} = \frac{\sqrt{2}}{2}(\mathbf{i} + \mathbf{j}) = \frac{\sqrt{2}}{2}\langle 1, 1 \rangle?$$

- | | | | |
|---|---|-----------------------------|---|
| <input type="checkbox"/> $2\sqrt{2}$ | <input type="checkbox"/> $\frac{\sqrt{2}}{4}$ | <input type="checkbox"/> 8 | <input type="checkbox"/> $\frac{\sqrt{6}}{2}$ |
| <input checked="" type="checkbox"/> $\frac{5\sqrt{2}}{2}$ | <input type="checkbox"/> $10\sqrt{2}$ | <input type="checkbox"/> -7 | <input type="checkbox"/> $\frac{1}{3}$ |

(d) (4 points). Which one of the following equations is an equation for the tangent plane to the graph of f at the point $Q = (1, 1, f(1, 1))$?

- | | |
|--|--|
| <input type="checkbox"/> $x + 2y - z = 11$ | <input checked="" type="checkbox"/> $11x + 4y + z = 7$ |
| <input type="checkbox"/> $2x - 3y + z = -9$ | <input type="checkbox"/> $x + y - z = 1$ |
| <input type="checkbox"/> $9x + 10y - z = 13$ | <input type="checkbox"/> $10x + y + z = 3$ |

Problem 9 (5 points)

Let

$$g(x, y, z) = \arctan(x^2 + y^2 - z^2).$$

In which direction is the directional derivative of g at $P = (1, -1, 1)$ maximal? Mark a unit vector below which specifies the direction.

- | | |
|---|--|
| <input checked="" type="checkbox"/> $\frac{\sqrt{3}}{3}\langle 1, -1, -1 \rangle$ | <input type="checkbox"/> $\frac{1}{3}\langle 2, 2, 1 \rangle$ |
| <input type="checkbox"/> $\frac{\sqrt{2}}{2}\langle 1, 1, 0 \rangle$ | <input type="checkbox"/> $\frac{1}{3}\langle -2, 1, 2 \rangle$ |
| <input type="checkbox"/> $\frac{\sqrt{2}}{2}\langle -1, 0, 1 \rangle$ | <input type="checkbox"/> $\frac{1}{5}\langle 3, -4, 0 \rangle$ |

Problem 10 (10 points)

A plane region \mathcal{R} consists of those points (x, y) which satisfy the inequalities

$$0 \leq x, \quad 0 \leq y, \quad x^2 + y^2 \leq 16.$$

Mark the value of the double integral

$$\iint_{\mathcal{R}} x \sqrt{x^2 + y^2} dA.$$

- | | | | | |
|--|---|-----------------------------|--|------------------------------|
| <input type="checkbox"/> $\frac{5}{4}$ | <input type="checkbox"/> $\frac{55}{4}$ | <input type="checkbox"/> 8 | <input checked="" type="checkbox"/> 64 | <input type="checkbox"/> 91 |
| <input type="checkbox"/> $\frac{5}{2}$ | <input type="checkbox"/> $\frac{47}{2}$ | <input type="checkbox"/> 55 | <input type="checkbox"/> 72 | <input type="checkbox"/> 121 |

Problem 11 (10 points)

A space region \mathcal{T} consists of those points (x, y, z) which satisfy the inequalities

$$0 \leq x \leq 1, \quad 0 \leq y \leq 2, \quad 0 \leq z \leq 5 - xy.$$

A solid body with density function $\delta(x, y, z) = x$ covers \mathcal{T} precisely.

(a) (5 points). What is the volume of \mathcal{T} ?

- | | | | | |
|--|--|---|----------------------------|---------------------------------------|
| <input type="checkbox"/> $\frac{1}{3}$ | <input type="checkbox"/> $\frac{3}{2}$ | <input type="checkbox"/> $\frac{22}{3}$ | <input type="checkbox"/> 3 | <input checked="" type="checkbox"/> 9 |
| <input type="checkbox"/> $\frac{1}{2}$ | <input type="checkbox"/> $\frac{7}{2}$ | <input type="checkbox"/> 2 | <input type="checkbox"/> 5 | <input type="checkbox"/> 11 |

(b) (5 points). What is the mass of the solid body covering \mathcal{T} ?


- | | | | | |
|--|---|----------------------------|----------------------------|---|
| <input checked="" type="checkbox"/> $\frac{13}{3}$ | <input type="checkbox"/> $\frac{15}{2}$ | <input type="checkbox"/> 3 | <input type="checkbox"/> 5 | <input type="checkbox"/> 10π |
| <input type="checkbox"/> $\frac{11}{2}$ | <input type="checkbox"/> 2 | <input type="checkbox"/> 4 | <input type="checkbox"/> 8 | <input type="checkbox"/> $\frac{4\pi}{7}$ |

Problem 12 (5 points)

The figure below shows the graph of the function

$$r = f(\theta), \quad 0 \leq \theta \leq 2\pi$$

in polar coordinates



Which one of the following rules for f corresponds to the figure?

$f(\theta) = 1 + \sin \theta$

$f(\theta) = \sin(2\theta)$

$f(\theta) = 1 + \theta$

$f(\theta) = 1 + \cos \theta$

$f(\theta) = 1 - 2 \sin \theta$

$f(\theta) = 1 + \theta^2$