

# Reexam in Calculus

First Year at the Faculty of Engineering and Science  
and the Technical Faculty of IT and Design

21 February 2018

The present exam set consists of 8 numbered pages with 12 multiple-choice problems. A number of points are specified for each question. The total number of points equals 100.

It is allowed to use books, notes etc. It **is not** allowed to use electronic devices.

Your answers must be marked directly in this exam set. The grading of the exam is based solely on these marks. Remember to write your **full name** and **student number** below.

Good luck!

NAME: \_\_\_\_\_

STUDENT NUMBER: \_\_\_\_\_

### Problem 1 (13 points)

(a) (5 points). A second order differential equation is given by

$$y'' - 10y' + 25y = 0.$$

Below is a list of function expressions containing two arbitrary constants  $c_1$  and  $c_2$ . Mark the expression which constitute the general solution of the differential equation.

- $y(t) = c_1e^{-3t} + c_2e^t$
- $y(t) = c_1e^{-5t} + c_2e^{2t}$
- $y(t) = c_1e^t + c_2e^{5t}$
- $y(t) = c_1e^{5t} + c_2te^{5t}$
- $y(t) = c_1e^{2t} + c_2te^{2t}$
- $y(t) = c_1e^{2t} \cos(5t) + c_2e^{2t} \sin(5t)$
- $y(t) = c_1e^t \cos(6t) + c_2e^t \sin(6t)$
- $y(t) = c_1e^{2t} \cos(3t) + c_2e^{2t} \sin(3t)$

(b) (4 points). Mark the solution to the initial value problem

$$y'' - 10y' + 25y = 0, \quad y(0) = 2, \quad y'(0) = 7$$

among the following options:

- $y(t) = -\frac{5}{4}e^{-3t} + \frac{13}{4}c_2e^t$
- $y(t) = 2e^{2t} + 3te^{2t}$
- $y(t) = 3e^{-5t} - c_2e^{2t}$
- $y(t) = 2e^{2t} \cos(5t) + 3e^{2t} \sin(5t)$
- $y(t) = -\frac{1}{4}e^t + \frac{5}{4}e^{5t}$
- $y(t) = 2e^t \cos(6t) + 4e^t \sin(6t)$
- $y(t) = 2e^{5t} - 3te^{5t}$
- $y(t) = e^{2t} \cos(3t) + e^{2t} \sin(3t)$

(c) (4 points). Consider the inhomogeneous differential equation

$$y'' - 10y' + 25y = 5t + 3.$$

Which one of the following functions is a particular solution to that equation?

- $\frac{5}{2}t + \frac{3}{2}$
- $t - 1$
- $t^2 - 5t - 2$
- $2e^t$
- $\frac{1}{5}t + \frac{1}{5}$
- $5t - 1$
- $2t^2 - t + 1$
- $3t - 4$

### Problem 2 (7 points)

A plane curve is given by

$$\begin{aligned}x &= t - e^t, \\y &= t + e^t,\end{aligned}$$

where the parameter  $t$  runs through the real numbers.

(a) (3 points). The point  $P = (-1, 1)$  lies on the curve. Which value of the parameter  $t$  corresponds to this point?

- |                                    |                               |                                   |  |
|------------------------------------|-------------------------------|-----------------------------------|--|
| <input type="checkbox"/> $-2$      | <input type="checkbox"/> $-1$ | <input type="checkbox"/> $1$      | <input type="checkbox"/> $3$           |
| <input type="checkbox"/> $-\ln(2)$ | <input type="checkbox"/> $0$  | <input type="checkbox"/> $\ln(2)$ | <input type="checkbox"/> $\frac{7}{2}$ |

(b) (4 points). What is the curvature of the curve for  $t = 0$ ?

- |   |   |  |  |
|---|---|--|--|
| <input type="checkbox"/> $\frac{5}{4}$        | <input type="checkbox"/> $\frac{\sqrt{3}}{4}$ | <input type="checkbox"/> $\frac{1}{5}$ | <input type="checkbox"/> $8$           |
| <input type="checkbox"/> $\frac{\sqrt{2}}{4}$ | <input type="checkbox"/> $\sqrt{5}$           | <input type="checkbox"/> $\frac{5}{3}$ | <input type="checkbox"/> $\frac{1}{4}$ |

### Problem 3 (8 points)

A curve in space is given by

$$\begin{aligned}x &= \frac{1}{2}t, \\y &= \frac{2}{3}t^{\frac{3}{2}}, \\z &= \frac{1}{2}t^2,\end{aligned}$$

where the parameter  $t$  runs through the positive real numbers.

(a) (3 points). What is the derivative  $y'$ ?

- |   |                                     |                                     |  |
|---|-------------------------------------|-------------------------------------|--|
| <input type="checkbox"/> $t^3 + 1$      | <input type="checkbox"/> $t \ln(t)$ | <input type="checkbox"/> $t$        | <input type="checkbox"/> $t\sqrt{t}$                   |
| <input type="checkbox"/> $\frac{1}{3t}$ | <input type="checkbox"/> $t^2$      | <input type="checkbox"/> $\sqrt{t}$ | <input type="checkbox"/> $\frac{4}{15}t^{\frac{5}{2}}$ |

(b) (5 points). What is the arc length of the curve from  $t = 1$  to  $t = 5$ ?

- |                               |                                       |                                      |   |
|-------------------------------|---------------------------------------|--------------------------------------|---|
| <input type="checkbox"/> $14$ | <input type="checkbox"/> $10\sqrt{2}$ | <input type="checkbox"/> $7\sqrt{3}$ | <input type="checkbox"/> $18$           |
| <input type="checkbox"/> $5$  | <input type="checkbox"/> $8$          | <input type="checkbox"/> $11$        | <input type="checkbox"/> $\frac{25}{2}$ |

### Problem 4 (7 points)

A function is defined by

$$f(x) = \frac{1}{x^2 + 2}.$$

(a) (3 points). What is the derivative  $f'(x)$ ?

- |  |   |  |
|--|---|--|
| <input type="checkbox"/> $\frac{1}{2x}$          | <input type="checkbox"/> $(x^2 + 2)^{-\frac{1}{2}}$ | <input type="checkbox"/> $2x(x^2 + 2)^{\frac{1}{2}}$ |
| <input type="checkbox"/> $-\frac{2x}{(x^2+2)^2}$ | <input type="checkbox"/> $-\frac{1}{(x^2+2)^2}$     | <input type="checkbox"/> $\ln(x^2 + 2)$              |

(b) (4 points). Which one of the polynomials below is the second order Taylor polynomial for  $f$  at the points  $x = 0$ ?

- |  |   |   |
|--|---|---|
| <input type="checkbox"/> $1 + x - \frac{3}{2}x^2$                      | <input type="checkbox"/> $\frac{1}{2} + x - \frac{1}{2}x^2$ | <input type="checkbox"/> $2 + 4x - 5x^2$      |
| <input type="checkbox"/> $\frac{1}{2} + \frac{1}{2}x + \frac{1}{6}x^2$ | <input type="checkbox"/> $\frac{1}{2} - \frac{1}{4}x^2$     | <input type="checkbox"/> $x - \frac{1}{2}x^2$ |

### Problem 5 (7 points)

Consider the differential equation

$$\frac{dy}{dx} = \frac{e^x}{y}, \quad y > 0.$$

There is a unique solution  $y(x)$  with initial value  $y(0) = 2$ . Answer the following questions regarding this solution:

(a) (3 points). What is the derivative  $y'(0)$ ?

- |   |  |                              |
|---|--|------------------------------|
| <input type="checkbox"/> $-2$           | <input type="checkbox"/> $\frac{1}{2}$ | <input type="checkbox"/> $2$ |
| <input type="checkbox"/> $-\frac{1}{2}$ | <input type="checkbox"/> $1$           | <input type="checkbox"/> $3$ |

(b) (4 points). What is the function value  $y(1)$ ?

- |                                  |  |  |
|----------------------------------|--|--|
| <input type="checkbox"/> $5$     | <input type="checkbox"/> $\frac{1}{3}$ | <input type="checkbox"/> $\ln(2)$        |
| <input type="checkbox"/> $e + 1$ | <input type="checkbox"/> $1$           | <input type="checkbox"/> $\sqrt{2e + 2}$ |

### Problem 6 (6 points)

Two complex numbers are given by

$$z_1 = \frac{4 + 3i}{2 - i} + 1 + i, \quad z_2 = (e^{2 + \frac{\pi}{6}i})^3$$

(a) (3 points). What is  $z_1$  written in standard form?

- |                                   |   |                                  |                                   |
|-----------------------------------|---|----------------------------------|-----------------------------------|
| <input type="checkbox"/> $2 + 3i$ | <input type="checkbox"/> $\frac{1}{4} + \frac{5}{4}i$ | <input type="checkbox"/> $1 - i$ | <input type="checkbox"/> $8i$     |
| <input type="checkbox"/> $1 + i$  | <input type="checkbox"/> $\frac{2}{3} + \frac{2}{3}i$ | <input type="checkbox"/> $-3i$   | <input type="checkbox"/> $-5 + i$ |

(b) (3 points). What is  $z_2$  written in standard form?

- |                                     |                                     |                                 |                                     |
|-------------------------------------|-------------------------------------|---------------------------------|-------------------------------------|
| <input type="checkbox"/> $1 + e^3i$ | <input type="checkbox"/> $1 - e^2i$ | <input type="checkbox"/> $-e^9$ | <input type="checkbox"/> $e + e^2i$ |
| <input type="checkbox"/> $e^6$      | <input type="checkbox"/> $1 + e^2i$ | <input type="checkbox"/> $e^6i$ | <input type="checkbox"/> $1 + 3i$   |

### Problem 7 (6 points)

A function is defined by

$$f(x, y) = x \cos(y) - y \sin(x).$$

(a) (2 points). What is the partial derivative  $f_x(x, y)$ ?

- |   |  |
|---|--|
| <input type="checkbox"/> $-\sin(y) - \cos(x)$ | <input type="checkbox"/> $\tan(y) + \sin(x)$   |
| <input type="checkbox"/> $\cos(y)$            | <input type="checkbox"/> $\cos(y) - y \cos(x)$ |
| <input type="checkbox"/> $\tan(y) - 1$        | <input type="checkbox"/> $x^2 \cos(y)$         |

(b) (4 points). The graph of  $f$  has a tangent plane at the point  $P = (\frac{\pi}{2}, 0, f(\frac{\pi}{2}, 0))$ . Mark an equation for this plane.

- |  |  |
|--|--|
| <input type="checkbox"/> $x - y - z = 0$             | <input type="checkbox"/> $x + 2y - 3z = \frac{\pi}{2}$ |
| <input type="checkbox"/> $2x + y - 2z = 0$           | <input type="checkbox"/> $x - y + z = \pi$             |
| <input type="checkbox"/> $x + y - z = \frac{\pi}{2}$ | <input type="checkbox"/> $x + y + 2z = -\pi$           |

### Problem 8 (16 points)

A function of two real variables is given by the transformation rule

$$f(x, y) = \frac{y}{1 + x^2 + y^2}.$$

Answer the questions below regarding this function.

(a) (4 points). The domain of  $f$  consists of all those points  $(x, y)$  which satisfy

- |  |   |
|--|---|
| <input type="checkbox"/> $x \geq 0$ and $y \geq 0$ | <input type="checkbox"/> $y \neq 0$                   |
| <input type="checkbox"/> $x > 0$ and $y > 0$       | <input type="checkbox"/> $x$ and $y$ are real numbers |
| <input type="checkbox"/> $x \neq 0$ and $y \neq 0$ | <input type="checkbox"/> $x \neq y$                   |

(b) (4 points). Which of the following points are critical points for  $f$ ? (*Remark: One false mark will cancel one true mark in this subquestion.*)

- |                                    |                                    |
|------------------------------------|------------------------------------|
| <input type="checkbox"/> $(0, 1)$  | <input type="checkbox"/> $(0, 2)$  |
| <input type="checkbox"/> $(1, 1)$  | <input type="checkbox"/> $(0, -1)$ |
| <input type="checkbox"/> $(1, -1)$ | <input type="checkbox"/> $(2, -1)$ |

(c) (4 points). What is the directional derivative  $D_{\mathbf{u}}f(P)$  at the point  $P = (1, 1)$  and in the direction of the unit vector  $\mathbf{u} = \frac{3}{5}\mathbf{i} + \frac{4}{5}\mathbf{j} = \langle \frac{3}{5}, \frac{4}{5} \rangle$ ?

- |   |  |
|---|--|
| <input type="checkbox"/> $\frac{8}{5}$  | <input type="checkbox"/> 1               |
| <input type="checkbox"/> $\frac{3}{25}$ | <input type="checkbox"/> $-\frac{3}{5}$  |
| <input type="checkbox"/> 2              | <input type="checkbox"/> $-\frac{2}{45}$ |

(d) (4 points). The level curve with equation  $f(x, y) = \frac{1}{4}$  can be described as:

- A parabola with equation  $y = x^2 + 3$ .
- A parabola with equation  $y = x^2 - x + 4$ .
- A straight line through  $(0, \sqrt{3})$  with slope 2.
- A straight line through  $(0, 3)$  with slope  $-\frac{1}{2}$ .
- A circle with center at  $(0, 2)$  and radius  $\sqrt{3}$ .
- A circle with center at  $(1, 1)$  and radius  $\frac{1}{3}$ .

**Problem 9 (7 points)**

A lamina covers a region  $\mathcal{R}$  in the plane precisely. The region  $\mathcal{R}$  consists of those points  $(x, y)$  which satisfy the inequalities

$$0 \leq x, \quad 0 \leq y, \quad 3x + y \leq 6.$$

The density of the lamina is  $\delta(x, y) = x$ . What is the mass of the lamina?

- |  |                            |  |                                      |                             |
|--|----------------------------|--|--------------------------------------|-----------------------------|
| <input type="checkbox"/> $\frac{1}{2}$ | <input type="checkbox"/> 2 | <input type="checkbox"/> $\frac{9}{2}$ | <input type="checkbox"/> $5\sqrt{2}$ | <input type="checkbox"/> 12 |
| <input type="checkbox"/> 1             | <input type="checkbox"/> 4 | <input type="checkbox"/> 5             | <input type="checkbox"/> 10          | <input type="checkbox"/> 14 |

**Problem 10 (10 points)**

A region  $\mathcal{R}$  in the plane consists of those points  $(x, y)$  which satisfy the inequalities

$$0 \leq x \leq y, \quad 1 \leq x^2 + y^2 \leq 9.$$

Mark the value of the double integral

$$\iint_{\mathcal{R}} \frac{y}{x^2 + y^2} dA.$$

- |  |                                     |  |                             |   |
|--|-------------------------------------|--|-----------------------------|---|
| <input type="checkbox"/> $\frac{1}{3}$ | <input type="checkbox"/> $\sqrt{2}$ | <input type="checkbox"/> $\frac{7}{2}$ | <input type="checkbox"/> 6  | <input type="checkbox"/> 12             |
| <input type="checkbox"/> $\frac{1}{2}$ | <input type="checkbox"/> 3          | <input type="checkbox"/> $5\sqrt{2}$   | <input type="checkbox"/> 10 | <input type="checkbox"/> $\frac{25}{2}$ |

**Problem 11 (8 points)**

A region  $\mathcal{T}$  in space consists of those points  $(x, y, z)$  which satisfy the inequalities

$$0 \leq x \leq 2, \quad 0 \leq y \leq 1, \quad 2y \leq z \leq 9 - 3xy^2.$$

What is the volume of  $\mathcal{T}$ ?


- |                            |  |                             |                             |   |
|----------------------------|--|-----------------------------|-----------------------------|---|
| <input type="checkbox"/> 2 | <input type="checkbox"/> $\frac{7}{2}$ | <input type="checkbox"/> 12 | <input type="checkbox"/> 16 | <input type="checkbox"/> $18\sqrt{3}$   |
| <input type="checkbox"/> 3 | <input type="checkbox"/> 9             | <input type="checkbox"/> 14 | <input type="checkbox"/> 20 | <input type="checkbox"/> $\frac{50}{3}$ |

**Problem 12 (5 points)**

The figure below shows the graph of the function

$$r = f(\theta), \quad 0 \leq \theta \leq 2\pi$$

in polar coordinates.



Which one of the following transformation rules for  $f$  corresponds to the figure?

$f(\theta) = 1 + \cos \theta$

$f(\theta) = 1 - \cos \theta$

$f(\theta) = 1 - \theta^2$

$f(\theta) = 8 \sin^2 \theta$

$f(\theta) = \theta(2\pi - \theta)$

$f(\theta) = 10 - \theta^2$