

Reexam in Calculus

19. august 2019

Exercise 1 (6 point)

A function is defined by

$$f(x, y, z) = \sin(x^2 + y^2 + z^2)$$

for real variables x , y and z .

- (a) (3 point) The domain of definition of f consists of all the points (x, y, z) which satisfy

- | | |
|--|---|
| <input type="checkbox"/> $z = 0$ | <input type="checkbox"/> all points are allowed |
| <input type="checkbox"/> $x^2 + y^2 + z^2 > 0$ | <input type="checkbox"/> $xyz \neq 0$ |
| <input type="checkbox"/> $x^2 + y^2 + z^2 < 0$ | <input type="checkbox"/> none of the others |

- (b) (3 point) Which points (x, y, z) belong to the level surface determined by $f(x, y, z) = 2$?

- A sphere given by $x^2 + y^2 + z^2 = \sin^{-1}(2)$
- The xy -plane
- There are no such points
- A plane parallel with the xy -plane given by $z = \sin^{-1}(2)$
- none of the others

Exercise 2 (6 point)

A parametric curve in space is given by

$$\mathbf{r}(t) = \langle \sin(t), \cos(t), t \rangle$$

where the parameter t is an arbitrary real number.

- (a) (2 point) What is the velocity vector of the curve?

- | | |
|---|---|
| <input type="checkbox"/> $\langle \sin(t), \cos(t), 1 \rangle$ | <input type="checkbox"/> $\langle \sin(t^2/2), \cos(t^2/2), t^2/2 \rangle$ |
| <input type="checkbox"/> $\langle \cos(t), \sin(t), 1 \rangle$ | <input type="checkbox"/> $\langle \sin(t^2/2), -\cos(t^2/2), t^2/2 \rangle$ |
| <input type="checkbox"/> $\langle \cos(t), -\sin(t), 1 \rangle$ | <input type="checkbox"/> none of the others |

- (b) (2 point) Which of the following vectors is the acceleration vector for $t = 2\pi$?

- $\langle 0, -1, 0 \rangle$ $\langle 1, 0, 0 \rangle$ $\langle 0, 1, 1/2 \rangle$
 $\langle 0, 1, 0 \rangle$ $\langle 1, 1, 1 \rangle$ none of the others

(c) (1 point) What is the speed?

- $\sqrt{\sin(t) + \cos(t) + t}$ 2 $\sqrt{t+1}$
 $\sqrt{1+t^2}$ $\sqrt{2}$ none of the others

(d) (1 point) What is the length of the curve from $t = 0$ to $t = 2$?

- 0 $2\sqrt{2}$ 1/2
 $\sqrt{2}$ $4\sqrt{2}$ none of the others

Exercise 3 (6 point)

Three complex numbers are given by

$$z_1 = 1 + 2i, \quad z_2 = 4 - 2i \quad \text{og} \quad z_3 = i^{100}.$$

(a) (2 point) What is $z_1 + z_2$ in polar form?

- 1 $5e^{-i\pi/2}$ 5
 $5e^{i\pi/2}$ $\sqrt{5}$ none of the others

(b) (2 point) What is $\frac{2z_1}{z_2}$ in the standard form $a + ib$?

- $1 + i$ $-1 - i$ 5
 i $-i$ none of the others

(c) (2 point) What is the principal argument of z_3 ?

- 0 $\pi/2$ π
 $\pi/4$ $3\pi/4$ none of the others

Hint to (c): The principal argument is a polar angle which belongs to the interval $] - \pi, \pi]$.

Exercise 4 (10 point)

(a) (5 point) A homogeneous second order differential equation is given by

$$y'' = -4y.$$

Below there are given a few functions where c_1 and c_2 are arbitrary real constants. Mark which expression represents the general solution to the differential equation.

$y(t) = c_1 e^{-2t} + c_2 e^{2t}$

$y(t) = c_1 e^t + c_2 t$

$y(t) = c_1 \cos(t) + c_2 \sin(t)$

$y(t) = c_1 \sin(2t) + c_2 \cos(2t)$

$y(t) = (c_1 + c_2 t) e^{-2t}$

$y(t) = (c_1 + c_2 t) e^{2t}$

$y(t) = c_1 + c_2$

none of the others

(b) (2 point) Which function $x_p(t)$ is a particular solution to the inhomogeneous differential equation

$$x''(t) = -4x(t) + 5e^t$$

among the following expressions:

$x_p(t) = e^{2t}$

$x_p(t) = e^{-t}$

$x_p(t) = e^{-2t}$

$x_p(t) = te^t$

$x_p(t) = e^t$

none of the others

(c) (3 point) Mark the solution $x(t)$ to the initial value problem

$$x''(t) = -4x(t) + 5e^t, \quad x(0) = 0, \quad x'(0) = 0,$$

among the following expressions:

$x(t) = \sin(t) - t \cos(t)$

$x(t) = -\sin(2t)/2 - \cos(2t) + e^t$

$x(t) = -\sin(t) + te^t$

$x(t) = t - te^{2t}$

$x(t) = \sin(2t)/2 + \cos(2t) - e^t$

none of the others

Exercise 5 (8 point)

Mark if the following statements are true or false:

(a) (2 point) $(\sqrt{2} + \sqrt{3})^2 = 5 + 2\sqrt{6}$.

True

False

(b) (2 point) $\left(\frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}}\right)^2 = i$.

True

False

(c) (2 point) If $f(x) = \sin(x)$ and $g(t) = \cos(t)$ then $h(t) = f(g(t))$ is differentiable and $h'(t) = -\sin(t) \cos(\cos(t))$.

True

False

(d) (2 point) $e^{-\ln(x)} = \frac{1}{x}$.

True

False

Opgave 6 (7 point)

A domain \mathcal{R} in the plane can be represented with the help of the inequality $x^2 + y^2 \leq 1$.

(a) (1 point) Which of the following curves describes the boundary of \mathcal{R} ?

a circle centred at $(0, 1)$ and with radius 1

a circle centred at $(0, 0)$ and with radius 1

a triangle with corners at the origin, $(0, 1)$ and $(1, 0)$

a square with side length 1 and center at the origin

none of the others

(b) (2 point) Which of the following inequalities characterize all the points of \mathcal{R} , where $(x, y) = (r \cos(\theta), r \sin(\theta))$?

$r \geq 1, \quad 0 \leq \theta \leq \pi$

$0 \leq r \leq 1, \quad 0 \leq \theta \leq 2\pi$

$0 \leq r \leq \cos(\theta), \quad 0 \leq \theta \leq \pi/2$

$r \geq 1$

$0 \leq r \leq \sin(\theta), \quad 0 \leq \theta \leq \pi$

none of the others

(c) (4 point) The domain \mathcal{R} models a plate with mass density $\delta(x, y) = x^2 + y^2$. Which of the following expressions give the mass of the plate?

- | | |
|---|--|
| <input type="checkbox"/> $2\pi \int_0^1 r^2 dr$ | <input type="checkbox"/> $\int_0^\pi \int_0^{\sin(\theta)} r^3 dr d\theta$ |
| <input type="checkbox"/> $2\pi \int_0^1 r^3 dr$ | <input type="checkbox"/> $\int_0^1 r^2 dr$ |
| <input type="checkbox"/> $\pi \int_0^1 r^3 dr$ | <input type="checkbox"/> none of the others |

Opgave 7 (8 point)

A domain \mathcal{R} in the plane consists of all the points with coordinates (x, y) which satisfy the inequality $x^2 + y^2 \leq 1$. The function f is defined on \mathcal{R} and given by $f(x, y) = x + y$.

(a) (4 point) Which of the following points are inner critical points for f ?

- | | |
|--|---|
| <input type="checkbox"/> $\langle 0, -1 \rangle$ | <input type="checkbox"/> $\langle -1, 0 \rangle$ |
| <input type="checkbox"/> $\langle 1, 0 \rangle$ | <input type="checkbox"/> There are no inner critical points |
| <input type="checkbox"/> $\langle 0, 0 \rangle$ | <input type="checkbox"/> none of the others |

(b) (2 point) Mark whether the following statement is true or false:

The function f evaluated on the boundary of \mathcal{R} takes the same values as the function $g(\theta) = \cos(\theta) + \sin(\theta)$, where $\theta \in [0, 2\pi]$.

- | | |
|-------------------------------|--------------------------------|
| <input type="checkbox"/> True | <input type="checkbox"/> False |
|-------------------------------|--------------------------------|

(c) (2 point) What is the maximal value of f ?

- | | | |
|-------------------------------------|--------------------------------------|---|
| <input type="checkbox"/> 1 | <input type="checkbox"/> 2 | <input type="checkbox"/> 3 |
| <input type="checkbox"/> $\sqrt{2}$ | <input type="checkbox"/> $2\sqrt{2}$ | <input type="checkbox"/> none of the others |

Exercise 8 (12 point)

A surface \mathcal{F} in space is determined by the equation $F(x, y, z) = 0$, where

$$F(x, y, z) = x^4 + y^4 - 2z^2$$

(a) (3 point) Which of the following expressions gives the gradient vector ∇F ?

- | | |
|--|---|
| <input type="checkbox"/> $\langle 4x, 4y, -4z \rangle$ | <input type="checkbox"/> $\langle x^3, y^3, -2 \rangle$ |
| <input type="checkbox"/> $\langle 4x^2, 4y^2, -4z^2 \rangle$ | <input type="checkbox"/> $\langle 0, 0, 0 \rangle$ |
| <input type="checkbox"/> $\langle 4x^3, 4y^3, -4z \rangle$ | <input type="checkbox"/> none of the others |

(b) (4 point) Which of the following equations represent the tangent plane to \mathcal{F} at the point $P = (1, 1, 1)$?

- | | | |
|--|---|---|
| <input type="checkbox"/> $0 = x + y + z$ | <input type="checkbox"/> $z = x + y$ | <input type="checkbox"/> $z = 2x - y$ |
| <input type="checkbox"/> $x + y - z = 1$ | <input type="checkbox"/> $z = y + 2x - 2$ | <input type="checkbox"/> none of the others |

(c) (5 point) From the equation $F(x, y, z) = 0$, what is the partial derivative $\partial z / \partial x$ at the point P ?

-1

-2

2

0

1

none of the others

Exercise 9 (12 point)

A function is given by

$$f(x, y) = \sin(2x + y),$$

where $x \geq 0$ and $y \geq 0$.

(a) (2 point) Mark whether the following statement is true or false: $f(x, y)$ can never be equal to zero.

True

False

(b) (2 point) Mark whether the following statement is true or false: $f(x, 0)$ is an increasing function of x .

True

False

(c) (4 point) What is the directional derivative $D_{\mathbf{u}}f(P)$ at the point $P = (0, 0)$ in the direction given by the unit vector $\mathbf{u} = \langle 0, 1 \rangle$?

0

3

4

1

2

none of the others

(d) (4 point) Which of the following unit vectors point in the direction in which f grows fastest at the point P (the direction \mathbf{v} for which $D_{\mathbf{v}}f(P)$ is largest)?

$\langle -\frac{2}{\sqrt{5}}, -\frac{1}{\sqrt{5}} \rangle$

$\langle \frac{2\sqrt{5}}{5}, -\frac{1}{\sqrt{5}} \rangle$

$\langle \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \rangle$

$\langle 1, 0 \rangle$

$\langle \frac{2\sqrt{5}}{5}, \frac{\sqrt{5}}{5} \rangle$

$\langle -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \rangle$

$\langle \frac{1}{\sqrt{5}}, \frac{2}{\sqrt{5}} \rangle$

$\langle 0, 1 \rangle$

none of the others

Exercise 10 (9 point)

A function is given by

$$f(x) = \cos(x^2)$$

for all real numbers x .

(a) (5 point) Mark the correct expression for $f''(x)$ (i.e. f twice differentiated)

- $-4 \cos(x^2)$
 $2 \cos(2x)$
 $-x^4 \cos(x^2)$
 $2 \sin(x^2) - x^2 \cos(x^2)$
 $-2 \sin(x^2) - 4x^2 \cos(x^2)$
 none of the others

(b) (4 point) Which of the following expressions represents the second order Taylor polynomial for f with the expansion point $x = 0$?

- $1 + x + x^2$
 $1 - x + x^2/2$
 $2x + x^2$
 $1 + x + x^2/2$
 1
 none of the others

Exercise 11 (11 point)

A curve in the plane is given by

$$\begin{aligned}
 x(t) &= t^2, \\
 y(t) &= \sin(t^3)
 \end{aligned}$$

for all real numbers t .

(a) (2 point) For which value of the parameter t does the curve go through the origin?

- π
 3π
 0
 2π
 4π
 none of the others

(b) (4 point) What is the value of the speed when $t = 0$?

- 0
 $\sqrt{2}$
 $\sqrt{5}$
 1
 $\sqrt{3}$
 none of the others

(c) (5 point) What is the acceleration vector at the origin?

- $\langle 0, 0 \rangle$
 $\langle 0, 1 \rangle$
 $\langle 2, 0 \rangle$
 $\langle 1, 0 \rangle$
 $\langle 1, 1 \rangle$
 none of the others

Exercise 12 (5 point)

Consider the following initial value problem

$$y'(x) = (x + 1)y(x), \quad y(0) = 1.$$

(a) (3 point) Assume that y solves the above equation and define

$$f(x) = \ln(y(x)).$$

Which initial value problem solves f ?

- | | |
|---|--|
| <input type="checkbox"/> $f'(x) = x + 1, f(0) = 1$ | <input type="checkbox"/> $f'(x) = e^x, f(0) = 0$ |
| <input type="checkbox"/> $f'(x) = -x - 1, f(0) = 1$ | <input type="checkbox"/> $f'(x) = 1/(x + 1), f(0) = 0$ |
| <input type="checkbox"/> $f'(x) = x + 1, f(0) = 0$ | <input type="checkbox"/> none of the others |

(b) (2 point) What is $y(x)$?

- | | | |
|------------------------------------|--|---|
| <input type="checkbox"/> $1 + x$ | <input type="checkbox"/> $1/(x + 1)$ | <input type="checkbox"/> $\ln(1 + x + x^2/2)$ |
| <input type="checkbox"/> $e^x - 1$ | <input type="checkbox"/> $e^{x+x^2/2}$ | <input type="checkbox"/> none of the others |