

Exam 2011

Mathematics for Multimedia Applikations
AAU-Cph, Medialogy

1. June 2011

Formalities

This exam set consists of 8 pages. There are 10 problems containing 34 sub-problems in total. Books and notes are allowed but *no electronic devices* such as calculators, computers or cell phones are permitted.

A number of points is indicated for every sub-problem. The sum of these points equals 100.

Date and time for the exam: 1. June, 9:00 - 13:00

You must indicate the following on each page:

- Full name
- Study number
- Page number

On the first page, you must indicate

- The total number of pages.

Remark that special values for sine and cosine are added as an appendix.

Good luck!

Problems

Problem 1.

- 1.a. (3 points) Find the derivative of the function $x^3 + 2x \sin(x)$ with respect to x .
- 1.b. (3 points) Let $f(x) = 4 \ln(x^3 + 2x)$. Calculate $f'(x)$.

Problem 2. Let $f(x) = x + 2 \cos(x)$. The graph of this function appears in figure 1.

- 2.a. (2 points) Compute $f'(x)$.
- 2.b. (3 points) Find an x such that $f'(x) = 0$.
- 2.c. (5 points) Describe all x such that $f'(x) = 0$.




Figure 1: The graph of the function $f(x) = x + 2 \cos(x)$ for $-10 \leq x \leq 20$.

Problem 3. Consider the graph of a function f in figure 2. Use the sheet on page 8 for your answers to the following:

3.a. (3 points) Indicate all points where $f'(x) = 0$.

3.b. (3 points) Sketch the graph of the derivative f' .




Figure 2: A function f .

Problem 4. Let $g(x) = e^{2x}$ where e is the base of the natural exponential function. Let


$$f(x) = \frac{\log_e(g(x)) - g'(x)}{2} + (e^x)^2.$$

4.a. (3 points) Find a reduced form of the function f which does not depend on the function g .

Problem 5. Let f be the function defined by

$$f(x) = \begin{cases} 1 - x^2, & x \in [-1, 0], \\ 1 - \sin(x), & x \in [0, \pi/2]. \end{cases}$$

The graph of f looks as follows:



5.a. (3 points) Compute $\int_{-1}^0 f(x) dx$.

5.b. (3 points) Compute $\int_0^{\pi/2} f(x) dx$.

5.c. (3 points) Find $\int_{-1}^{\pi/2} f(x) dx$.

Problem 6. Let P , Q and R be three points in 3D-space; P has coordinates $(1, 2, 2)$, Q has coordinates $(1, 5, 2)$ and R has coordinates $(0, 3, 2)$.

6.a. (2 points) Find \overrightarrow{PQ} and \overrightarrow{PR} .

6.b. (3 points) Write parametric equations of the line L_1 that passes through P and Q and the line L_2 that passes through P and R .

6.c. (4 points) The lines L_1 and L_2 intersect at the point P . Find the angle between the two lines.


Problem 7. Let P , Q and R be three points in 3D-space; P has coordinates $(2, 1, 5)$, Q has coordinates $(3, 0, 5)$ and R has coordinates $(3, 1, 7)$.

- 7.a. (2 points) Find \overrightarrow{PQ} and \overrightarrow{PR} .
- 7.b. (3 points) Compute the cross product $\overrightarrow{PQ} \times \overrightarrow{PR}$.
- 7.c. (3 points) Find the area of the triangle with vertices P , Q and R .
- 7.d. (3 points) Find an equation for the plane through P , Q and R .
- 7.e. (2 points) Does the point $(3, -3, -1)$ lie on the plane? Why/why not?

Problem 8. A parametric curve is given by the following vector function:

$$\vec{r}(t) = (-\cos(t\sqrt{2}), \sin(t\sqrt{2}), t)$$

Here is a plot of the curve when t runs from 0 to 12 :



- 8.a. (3 points) Show that every point on the curve lies on the cylinder described by $x^2 + y^2 = 1$.
- 8.b. (2 points) Compute $\vec{r}'(t)$.
- 8.c. (3 points) Find a t between 0 and 2π such that the velocity vector of the parametric curve equals $(1, 1, 1)$.

Problem 9. Consider the following system of linear equations:

$$\begin{aligned}x_1 - 4x_3 &= 4 \\3x_2 - 2x_3 &= 3 \\-2x_1 + 6x_2 + 4x_3 &= -2\end{aligned}$$

- 9.a. (2 points) Find the augmented matrix of the system.
- 9.b. (4 points) Find a row echelon form of the augmented matrix.
- 9.c. (3 points) Find the reduced row echelon form of the augmented matrix.
- 9.d. (4 points) Write down the general solution to the system.
- 9.e. (3 points) One solution to the system is $x_1 = 4$, $x_2 = 1$, $x_3 = 0$. Find another solution which has $x_3 = 3$.
- 9.f. (3 points) Find a solution to the system which has $x_1 = -8$.

Problem 10. Define four matrices as follows:

$$A = \begin{bmatrix} 4 & 3 \\ 7 & -8 \\ -6 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 7 & -3 \\ 2 & 1 \end{bmatrix}, \quad C = \begin{bmatrix} 0 & 3 \\ 1 & 1 \\ 2 & 2 \end{bmatrix}, \quad D = \begin{bmatrix} 2 & 4 & 3 \\ 0 & 1 & 0 \\ 1 & 3 & 1 \end{bmatrix}$$

- 10.a. (2 points) Find A^T .
- 10.b. (3 points) Compute $A^T C$ or $C A^T$.
- 10.c. (3 points) Compute $(C^T A)^T + B$.
- 10.d. (4 points) Find D^{-1} .
- 10.e. (2 points) Let $T(\vec{x}) = D\vec{x}$. Compute $T\left(\begin{bmatrix} 3 \\ 0 \\ -1 \end{bmatrix}\right)$.
- 10.f. (3 points) Find an \vec{x} such that $T(\vec{x}) = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$.

Appendix

Exact values of sin and cos for some angles:

- $\sin(\pi/6) = \cos(\pi/3) = 1/2$
- $\sin(\pi/4) = \cos(\pi/4) = 1/\sqrt{2} = \sqrt{2}/2$
- $\sin(\pi/3) = \cos(\pi/6) = \sqrt{3}/2$




Figure 3: The function f